



Hyperon Photoproduction: What Has Been Learned at Jefferson Lab?



Reinhard Schumacher
Carnegie Mellon University

for the CLAS & GlueX Collaborations

May 20, 2015, CIPANP, Vail, Colorado




Outline / Overview

- Strangeness and the N^* spectrum of states
 - Λ & Σ photo- and electro-production spin observables
- Dimensional scaling of $K\Lambda$ photoproduction
 - Constituent-counting rule supported
- Excited Y^* cross sections measured at CLAS
 - $\Sigma^0(1385)$ ($J^P = 3/2^+$); $\Lambda(1405)$ ($J^P = 1/2^-$); $\Lambda(1520)$ ($J^P = 3/2^-$)
- Structure of the $\Lambda(1405)$: $\Sigma \pi$ line shapes; J^P
 - Support for chiral unitary models: 2-pole structure
- Strangeness suppression in exclusive electro-production
 - Low and high energy reactions similar behavior
- Outlook at GlueX and CLAS12

γ Y^*
 K^+ | Jefferson Lab


- Located in Newport News, Virginia
- Ran for ~14 yrs at 6 GeV in Halls A, B, C
- Upgrading to 12 GeV, new Hall D



- Most Y , Y^* publications from Hall B
 - Upgrading as CLAS12 for 12 GeV



CLAS Experiment




CLAS Experiment

■ Photoproduction:


- Targets: unpolarized LH_2 , polarized p, & HD-ice
- Beams: unpolarized, circular, linear, to $\sim 5 \text{ GeV}$
- Reconstructed $\text{K}^+p\pi^-(\pi^0)$ or $\text{K}^+\pi^+\pi^-(n)$
- 20×10^9 triggers $\rightarrow 1.41 \times 10^6$ $\text{KY}\pi$ events in g11a

■ Electroproduction:

- Q^2 from ~ 0.5 to $\sim 3 (\text{GeV}/c)^2$
- Structure functions from Rosenbluth and beam-helicity separations



Strangeness and the N^* Spectrum of States - Photoproduction



Strangeness in N^* Physics: Status

Table 8. Star rating suggested for baryon resonances and their decays. Ratings of the Particle Data Group are given as *; additional stars suggested from this analysis are represented by *; (*) stands for stars which should be removed.

	all	πN	γN	$N\eta$	ΔK	ΣK	$\Delta\pi$	$N\sigma$
$N(1440)_{1/2}^+$	****	****	****	(*)			***	***
$N(1710)_{1/2}^+$	***	***	***	***	***	***	*(*)	
$N(1880)_{1/2}^+$	**	*	*		**	*		
$N(1535)_{1/2}^-$	****	****	****	****			*	
$N(1650)_{1/2}^-$	****	****	***	***	***	**	**(*)	
$N(1895)_{1/2}^-$	**	*	**	**	**	*		
$N(1720)_{1/2}^+$	****	****	****	****	**	**	***	
$N(1900)_{1/2}^+$	***	**	***	**	***	***	**	
$N(1520)_{3/2}^-$	****	****	****	***			****	
$N(1700)_{3/2}^-$	**	**	**	*	*(*)	*	***	
$N(1875)_{3/2}^-$	***	*	***		***	***		***
$N(2150)_{3/2}^-$	**	**	**		**		**	
$N(1680)_{3/2}^+$	****	****	****	*			**(*)	**
$N(1860)_{3/2}^+$	*	*	*					
$N(2000)_{3/2}^+$	***	*(*)	**	**	**	*		
$N(1675)_{3/2}^-$	****	****	****(*)	*	*		**(*)	*
$N(2060)_{3/2}^-$	***	**	***	*		**		
$N(1990)_{7/2}^+$	**	*(*)	**					
$N(2190)_{7/2}^-$	****	****	***		**			
$N(2220)_{9/2}^+$	****	****						
$N(2250)_{9/2}^-$	****	****						
$\Delta(1910)_{1/2}^+$	****	****	**		**		**	
$\Delta(1620)_{1/2}^-$	****	****	**				****	
$\Delta(1900)_{1/2}^-$	**	**	**		**		**	
$\Delta(1232)_{3/2}^+$	****	****	****					
$\Delta(1600)_{3/2}^+$	***	**	**				***	
$\Delta(1920)_{3/2}^+$	***	**	**		***		**	
$\Delta(1700)_{3/2}^-$	***	**	**				**	
$\Delta(1940)_{3/2}^-$	*	*	**					* from $\Delta\eta$
$\Delta(1905)_{5/2}^+$	****	****	****		***		**(*)	
$\Delta(1950)_{7/2}^+$	****	****	**		***		**	



S_{11} →
 P_{13} →
 D_{13} →
 G_{17} →

- What role has JLab strangeness physics in unraveling N^* and Δ properties?
- Worldwide effort to determine resonance poles, branching fractions, helicity couplings, etc.
- Bottom line: "Stars" & new resonances added to world database


γ Y^*
 K^+

Define the Spin Observables

(for unpolarized nucleon)



$$\frac{d\sigma}{d\Omega} = \sigma_0 \left\{ \begin{array}{l} 1 - P_{\vec{\gamma}} \Sigma \cos 2\phi \\ -\alpha \cos \theta_{x'} \sin 2\phi P_{\vec{\gamma}} O_{x'} - \alpha \cos \theta_{x'} P_{\odot} C_{x'} \\ -\alpha \cos \theta_{z'} \sin 2\phi P_{\vec{\gamma}} O_{z'} - \alpha \cos \theta_{z'} P_{\odot} C_{z'} \\ + \alpha \cos \theta_{y'} P - \alpha \cos \theta_{y'} P_{\vec{\gamma}} T \cos 2\phi \end{array} \right\}$$




Pseudoscalar Meson Photoproduction

4 Complex amplitudes: **16** real polarization observables.

Complete measurement with at least **8** suitably chosen observables.

nN has large cross section

but in **KY** recoil is **self-analysing**



nN		Symbol	Transversity representation	Experiment required	Type	KY		
recoil	target	γ				γ	target	recoil
		$d\sigma/dt$	$ b_1 ^2 + b_2 ^2 + b_3 ^2 + b_4 ^2$	$\{-; -; -\}$	<i>S</i>			
		$\Sigma d\sigma/dt$	$ b_1 ^2 + b_2 ^2 - b_3 ^2 - b_4 ^2$	$\{L(\frac{1}{2}\pi, 0); -; -\}$				
		$Td\sigma/dt$	$ b_1 ^2 - b_2 ^2 - b_3 ^2 + b_4 ^2$	$\{-; y; -\}$				
		$Pd\sigma/dt$	$ b_1 ^2 - b_2 ^2 + b_3 ^2 - b_4 ^2$	$\{-; -; y\}$				
		$Gd\sigma/dt$	$2 \text{Im}(b_1 b_3^* + b_2 b_4^*)$	$\{L(\pm\frac{1}{4}\pi); z; -\}$	<i>BT</i>			
		$Hd\sigma/dt$	$-2 \text{Re}(b_1 b_3^* - b_2 b_4^*)$	$\{L(\pm\frac{1}{4}\pi); x; -\}$				
		$Ed\sigma/dt$	$-2 \text{Re}(b_1 b_3^* + b_2 b_4^*)$	$\{C; z; -\}$				
		$Fd\sigma/dt$	$2 \text{Im}(b_1 b_3^* - b_2 b_4^*)$	$\{C; x; -\}$				
		$O_x d\sigma/dt$	$-2 \text{Re}(b_1 b_4^* - b_2 b_3^*)$	$\{L(\pm\frac{1}{4}\pi); -; x'\}$	<i>BR</i>			
		$O_z d\sigma/dt$	$-2 \text{Im}(b_1 b_4^* + b_2 b_3^*)$	$\{L(\pm\frac{1}{4}\pi); -; z'\}$				
		$C_x d\sigma/dt$	$2 \text{Im}(b_1 b_4^* - b_2 b_3^*)$	$\{C; -; x'\}$				
		$C_z d\sigma/dt$	$-2 \text{Re}(b_1 b_4^* + b_2 b_3^*)$	$\{C; -; z'\}$				
		$T_x d\sigma/dt$	$2 \text{Re}(b_1 b_2^* - b_3 b_4^*)$	$\{-; x; x'\}$	<i>TR</i>			
		$T_z d\sigma/dt$	$2 \text{Im}(b_1 b_2^* - b_3 b_4^*)$	$\{-; x; z'\}$				
		$L_x d\sigma/dt$	$2 \text{Im}(b_1 b_2^* + b_3 b_4^*)$	$\{-; z; x'\}$				
		$L_z d\sigma/dt$	$2 \text{Re}(b_1 b_2^* + b_3 b_4^*)$	$\{-; z; z'\}$				

I. S. Barker, A. Donnachie, J. K. Storrow, Nucl. Phys. B95 347 (1975).

circ polarized photons
 linearly polarized photons

longitudinally polarized target
 transversely polarized target


γ Y^* K^+ Theory: Bonn Gatchina Model

(Just one of several models on the market)

- Coupled channels (K-matrix) framework
 - Input: from πN , $K N$ elastic; γN , πN inelastic to $\pi^{\pm 0} N$, ηN , $\eta' N$, $K^{\pm 0} Y$, $\pi \pi N$
 - Use ALL experimental channels, including the strangeness channels & spin observables
 - Partial Wave Analysis
 - First extract each J^P wave
 - Fit N^* and Δ resonance pole parameters


Short list of References:

- A. Sarantsev, V. Nikonov, A. Anisovich, E. Klempt, U. Thoma; Eur. Phys. J. A **25**, 441 (2005)
A.V. Anisovich *et al.*, Eur. Phys J. A **25** 427 (2005); Eur. Phys J. A **24**, 111 (2005);
V. A. Nikonov *et al.*, Phys Lett. B **662**, 246 (2008).
A. Anisovich, E. Klempt, V. Nikonov, A. Sarantsev, U. Thoma; Eur. Phys. J. A **47**, 153 (2011).







$\gamma p \rightarrow K^+ \Lambda$: cross section

R. Bradford *et al.*, Phys.Rev. C **73**, 035202 (2006)




- Forward peaking indicates t-channel processes at high W
- Angular dependence at lower W consistent with s- and u-channel processes.

$\gamma Y^* K^+ | \gamma p \rightarrow K^+ \Lambda$: recoil polarization P



- Kaon-MAID model (green)
 - F.X. Lee *et al.*, Nucl. Phys. **A695**, 237 (2001).
 - Single-channel BW resonance fits
 - No longer up-to-date
- Bonn-Gatchina model (blue)
 - Multi-channel, unitary, BW resonance fit
 - Large suite of N^* contributions
 - Was not predictive for recoil polarization


A.V. Sarantsev *et al.*, Eur. Phys. J., A **25**, 441 (2005).



$\gamma p \rightarrow K^+ \Lambda$: beam asymmetry Σ

$$\frac{d\sigma}{d\Omega_{K^+}} = \frac{d\sigma}{d\Omega_{K^+}} \Big|_{unpol.} \{1 + \Sigma P_\gamma \cos 2\phi\}$$


Observable: Beam Asymmetry, Σ



Bonn-Gatchina model is not predictive in newly-measured kinematics

γ K^+ Y^* | $\gamma p \rightarrow \text{K}^+ \Lambda$: target asymmetry T


$$\gamma + p \rightarrow \text{K}^+ \Lambda$$



Bonn-Gatchina model is not predictive in newly-measured kinematics


γ γ^* K^+ $\vec{\gamma}p \rightarrow K^+ \vec{\Lambda}$ Beam-Recoil O_x and O_z

$\gamma + p \rightarrow K^+ \Lambda$




O_x

The Bonn-Gatchina model is not predictive at newly-measured kinematics




O_z

γ Y^* K^+ $\vec{\gamma}p \rightarrow K^+ \vec{\Lambda}$ Beam-Recoil C_x and C_z




$C_x C_z$ without $N^*(1900)P_{13}$



$C_x C_z$ with $N^*(1900)P_{13}$


- Nikanov *et al.*'s refit of Bonn-Gatchina coupled-channel isobar model
- mix includes: S_{11} -wave, $P_{13}(1720)$, $P_{13}(1900)$, $P_{11}(1840)$
- $K^+\Sigma^0$ cross sections also better described with $P_{13}(1900)$



R. Bradford *et al.*, (CLAS Collaboration) Phys. Rev. C **75**, 035205 (2007).


V. A. Nikanov *et al.*, Phys Lett. B **662**, 246 (2008).

see also: A.V. Anisovich *et al.*, Eur. Phys J. A **25** 427 (2005).




Seeking New $S=0$ Baryons via Mesons off the Proton: published, acquired, FroST(g9b)

	σ	Σ	T	P	E	F	G	H	T_x	T_z	L_x	L_z	O_x	O_z	C_x	C_z	CLAS run Period
$p\pi^0$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g8, g9
$n\pi^+$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g8, g9
$p\eta$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g11, g8, g9
$p\eta'$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g11, g8, g9
$p\omega$	✓	✓	✓	✓	✓	✓	✓	✓									g11, g8, g9
$K^+\Lambda$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	g1, g8, g11
$K^+\Sigma^0$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	g1, g8, g11
$K^{0*}\Sigma^+$	✓										✓	✓			✓	✓	g1, g8, g11



Lots more could be said...


- Omit results for Σ photoproduction
- Omit discussion of reactions on the neutron (deuteron), which accesses photon coupling isospin dependence.
- Overall goal: measure enough observables for "complete" determination of amplitudes \Rightarrow extract N^* and Δ content



Strangeness and the N^* Spectrum of States - Electroproduction

Structure Functions

For unpolarized target & polarized e^- beam:



$$\frac{d^4\sigma}{dQ^2 dW d\Omega_K} = \Gamma(Q^2, W) \times \frac{d\sigma}{d\Omega_K}(Q^2, W, \Theta_K, \varepsilon, \phi)$$

Virtual
photon
flux

Meson cross section

Transverse

Transverse-transverse
interference

Helicity
structure


$$\frac{d\sigma}{d\Omega_K} = \sigma_T + \varepsilon_L \sigma_L + \varepsilon \sigma_{TT} \cos(2\phi) + \sqrt{2\varepsilon_L(\varepsilon+1)} \sigma_{LT} \cos(\phi) + h\sqrt{2\varepsilon_L(1-\varepsilon)} \sigma_{LT'}$$

σ_u
"Unseparated"

Longitudinal (sensitive
to $J=0^\pm$ exchange in
t-channel: kaons, diquarks)


Transverse-longitudinal
interference

$\gamma Y^* | K^+ \Lambda$ Structure Functions



$E = 5.5 \text{ GeV}$, $W: \text{thr} - 2.6 \text{ GeV}$, $Q^2 = 1.80, 2.60, 3.45 \text{ GeV}^2$ [Carman *et al.*, PR C **87**, 025204 (2013)]

$\gamma Y^* K^+$ | L/T Separation at $Q^2=1.0 \text{ (GeV/c)}^2$



[Ambrozewicz *et al.*, PR C **75**, 045203 (2007)]


γ Y^* K^+ CLAS $e p$ Data Set Overview

#	Period	E_b (GeV)	Events (M)
1	e1c	2.567	900
2	e1c	4.056	370
3	e1c	4.247	620
4	e1c	4.462	420
5	e1d	4.817	300
6	e1-6	5.754	4500
7	e1f	5.499	5000
8	e1g	3.178	2500

- $K^+\Lambda$ recoil polarization
 - $W=1.6-2.7$ GeV, $\langle Q^2 \rangle = 1.9$ GeV²
[Gabrielyan *et al.*, PR C **90**, 035202 (2014)]

Publications:

- $K^+\Lambda$ beam-recoil pol. transfer
 - $W=1.6-2.15$ GeV, $Q^2=0.3 - 1.5$ GeV²
[Carman *et al.*, PRL **90**, 131804 (2003)]
- $K^+\Lambda$ σ_L/σ_T ratio from pol. transfer data
 - $W=1.72-1.98$ GeV, $Q^2 \sim 0.7$ GeV²
[Raue & Carman, PR C **71**, 065209 (2005)]
- $K^+\Lambda$, $K^+\Sigma^0$ separated structure functions
 - $W=thr-2.4$ GeV, $Q^2=0.5-2.8$ GeV²
 - $\sigma_U, \sigma_{LT}, \sigma_{TT}, \sigma_L, \sigma_T$ - $K^+\Lambda, K^+\Sigma^0$
[Ambrozewicz *et al.*, PR C **75**, 045203 (2007)]
 - $W=thr-2.6$ GeV, $Q^2=1.4-3.9$ GeV²
 - $\sigma_U, \sigma_{LT}, \sigma_{TT}, \sigma_{LT}$ - $K^+\Lambda, K^+\Sigma^0$
[Carman *et al.*, PRC **87**, 025204 (2013)]
- $K^+\Lambda$ fifth structure function σ_{LT}
 - $W=1.6-2.1$ GeV, $Q^2=0.65, 1.0$ GeV²
[Nasseripour *et al.*, PR C **77**, 065208 (2008)]
- $K^+\Lambda, K^+\Sigma^0$ beam-recoil pol. transfer
 - $W=thr-2.6$ GeV, $Q^2=1.6-2.6$ GeV²
[Carman *et al.*, PR C **79**, 065205 (2009)]





Dimensional Scaling of $K\Lambda$


Publication: **Scaling and Resonances in Elementary $K^+\Lambda$ Photo-production**, R.A.Sch. and M.M. Sargsian Phys.Rev.C**83** 025207 (2011).

Constituent-Counting Scaling


$$\frac{d\sigma}{dt} = f\left(\frac{t}{s}\right) s^{2-n}$$




- Constituent counting rules for exclusive scattering
- Valid for $s \rightarrow \infty$ and t/s fixed
 - $t/s \sim \cos(\theta_{\text{cm}})$ as $s \rightarrow \infty$
- n = number of point-like constituents
- Follows from pQCD... but also other models
- Does it work for $K\Lambda$?




Resonance Fit to Cross Section






Excited Y^* Cross Sections

Publication: **Differential Photoproduction Cross Sections of $\Sigma^0(1385)$, $\Lambda(1405)$ and $\Lambda(1520)$** , K. Moriya *et al.* (CLAS Collaboration), Phys. Rev. C **88**, 045201 (2013).




Detect $K^+p\pi^-(\pi^0)$ or $K^+\pi^+\pi^-(n)$

counts/5 MeV $MM(\gamma, K^+)$



γ Y^* K^+ Differential $\Sigma^0(1385)$ Cross Section



- $\gamma + p \rightarrow K^+ + \Sigma^0(1385)$

- Experiment: see t -channel-like forward peaking & u -channel backward rise


- Agreement with LEPS

- Theory by Oh et al.¹: contact term dominant; included four high-mass N^* and Δ resonances

- Prediction was fitted to preliminary CLAS total cross section (years ago)

1. Y. Oh, C. M. Ko, K. Nakayama, Phys. Rev. **C 77**, 045204 (2008)

γ Y^* K^+ Differential $\Lambda(1520)$ Cross Section



■ $\gamma + p \rightarrow K^+ + \Lambda(1520)$

■ Experiment: see t -channel-like forward peaking & u -channel backward rise

■ Agreement with LEPS^{1,2}


■ Theories:

■ Nam & Kao³: contact term dominant; no K^* or u -channel exchanges


■ He & Chen⁴: K^* and $N(2080)D_{13}$ $J^P=3/2^-$ added

1. H. Kohri et al. (LEPS) Phys Rev Lett **104**, 172001 (2010)
2. N. Muramatsu et al. (LEPS) Phys Rev **103**, 012001 (2009)
3. S.I. Nam & C.W. Kao, Phys. Rev. **C 81**, 055206 (2010)
4. J. He & X.R. Chen, Phys. Rev. **C 86**, 035204 (2012)


γ Y^* K^+ Differential $\Lambda(1405)$ Cross Section



- $\gamma + p \rightarrow K^+ + \Lambda(1405)$
- Experiment: first-ever measurements
- Low W: See strong isospin dependence
 - Charge channels differ
 - WHY?!?
- High W: See t -channel-like forward peaking & u -channel backward rise at high W
- Channels merge together at high W




Total Cross Sections Comparison



- $\gamma + p \rightarrow K^+ + Y^*$
- All three Y^* s have similar total cross sections
- Ground state Λ and Σ^0 are comparable to Y^* in size¹


1. R. Bradford et al. (CLAS) Phys. Rev. **C 73**, 035202 (2006)
K. Moriya *et al.* (CLAS), Phys. Rev. C **88**, 045201 (2013).



$\Lambda(1405)$ Structure


Publications: Measurement of the $\Sigma\pi$ Photo-production Line Shapes Near the $\Lambda(1405)$, K. Moriya *et al.* (CLAS Collaboration), Phys. Rev. C **87**, 035206 (2013);


Isospin Decomposition of the Photoproduced $\Sigma\pi$ System near the $\Lambda(1405)$, R. A. Sch. & K. Moriya, Nucl. Phys A **914**, 51 (2013) .



What "is" the $\Lambda(1405)$?

- An issue since its prediction/discovery
 - Dynamically generated resonance, via unitary meson-baryon channel coupling
 - R. Dalitz & S.F. Tuan, Phys. Rev. Lett. 2, 425 (1959), Ann. Phys. 10, 307 (1960).
 - Chiral unitary models (present-day theoretical industry!)
 - $SU(3)$ singlet $3q$ state, $I=0$, $J^\pi = \frac{1}{2}^-$
- $\bar{K}N$ sub-threshold state
 - Recent first Lattice QCD result:
 - J. Hall *et al.*, Phys Rev Lett 114, 132002 (2015)
- Signal may be an overlay of $I=0$ and $I=1$ states





Chiral Unitary Models






Fig. 1. Trajectories of the poles in the scattering amplitudes obtained by changing the SU(3) breaking parameter x gradually. At the SU(3) symmetric limit ($x = 0$), only two poles appear, one is for the singlet and the other for the octets. The symbols correspond to the step size $\delta x = 0.1$.

- SU(3) baryons irreps $1+8_s+8_a$ combine with 0- Goldstone bosons to generate:
- Two octets and a singlet of $\frac{1}{2}^-$ baryons generated dynamically in SU(3) limit
- SU(3) breaking leads to two $S = -1$ $I = 0$ poles near 1405 MeV
 - ~1420 mostly $\bar{K}N$
 - ~1390 mostly $\pi\Sigma$
- Possible weak $I=1$ pole also predicted

D. Jido, J.A. Oller, E. Oset, A. Ramos, U-G Meissner Nucl. Phys. A **725**, 181 (2003)
 J.A. Oller, U.-G. Meissner Phys. Lett B **500**, 263 (2001).




Example at $W=2.30$ GeV



I=1 contribution

Isospin Interference



$$|t_I|^2 \equiv |\langle I, 0 | \hat{T}^{(I)} | \gamma p \rangle|^2$$



Three charge combinations:

$$|T_{\pi-\Sigma^+}|^2 = \frac{1}{3}|t_0|^2 + \frac{1}{2}|t_1|^2 - \frac{2}{\sqrt{6}}|t_0||t_1|\cos\phi_{01},$$



$$|T_{\pi^0\Sigma^0}|^2 = \frac{1}{3}|t_0|^2,$$

$$|T_{\pi^+\Sigma^-}|^2 = \frac{1}{3}|t_0|^2 + \frac{1}{2}|t_1|^2 + \frac{2}{\sqrt{6}}|t_0||t_1|\cos\phi_{01}.$$

γ




Line Shapes



Cross Sections

- Charge-dependence is NOT seen for the $\Lambda(1520)$.
- No model calculation has computed cross section and line shapes together.




What "is" the I=1 piece?

- I=1 resonance? I=1 continuum amplitude?
- L. Roca and E. Oset model¹:
 - Possible I=1 resonance in vicinity of $N\bar{K}$ threshold
- B.-S. Zou et al. model²:
 - $\Sigma\left(\frac{1}{2}\right)^-$ is a $|[ud][us]\bar{s}\rangle$ state: part of a new nonet
- No interference seen in $\Lambda(1520)$ mass range: therefore it's not a continuum amplitude
- More investigation needed !

1. L. Roca, E. Oset "On the isospin 0 and 1 resonances from $\pi\Sigma$ photoproduction data" Phys. Rev. C **88** 055206 (2013).

2. Bing-Song Zou "Five-quark components in baryons", Nucl Phys A 835 199 (2010).



Spin and Parity of $\Lambda(1405)$

Publication: **Spin and Parity of the $\Lambda(1405)$ Baryon**, K. Moriya *et al.*
(CLAS Collaboration), Phys. Rev. Lett. **112**, 082004 (2014).

γ Y^* K^+ | Parity and Spin of $\Lambda(1405)$

- How does one measure these things?
 - Find a reaction wherein Λ^* is created polarized
 - Decay angular distribution to $\Sigma \pi$ relates to J
 - $J = 1/2$: flat distribution is the best possible evidence
 - $J = 3/2$: "smile or frown" distribution, where p is the $m = \pm 3/2$ fraction

$$I(\theta_Y) \propto 1 + \frac{3(1-2p)}{2p+1} \cos^2 \theta_Y$$


- Parity given by polarization transfer to daughter
- No model dependence: pure kinematics

γ Y^*
 K^+

S-wave, P-wave Scenarios


L=0 (s-wave)

$$\vec{Q} = \vec{P}$$




L=1 (p-wave)

$$\vec{Q} = -\vec{P} + 2(\vec{P} \cdot \hat{q})\hat{q}$$



$\Lambda(1405) \rightarrow \Sigma \pi$


γ Y^* K^+ Parity and Spin of $\Lambda(1405)$



$J^P = 1/2^-$ confirms quark model expectation

- Polarization axis is along $\hat{z} = \hat{\gamma} \times \hat{K}$
- Used $W=2.55$ to 2.85 GeV, $\cos \theta_K^{c.m.} > 0.6$
- Decay $\Lambda(1405) \rightarrow \Sigma^+ \pi^-$ is isotropic ($p = 0.5$), so $J \rightarrow 1/2$
- Weak decay asymmetry for Σ^+ is $\alpha = -0.98$ (big!)
- Decay is s-wave, $\Rightarrow P = \text{"negative"}$




and $\Lambda(1405)$ is produced $\sim +45\%$ polarized




$\Lambda(1405)$ Electroproduction

Publication: **First Observation of the $\Lambda(1405)$ Line Shape in Electroproduction**, H. Lu *et al.* (CLAS Collaboration), *Phys. Rev. C* **88**, 045202 (2013).

γ Y^* K^+ Electroproduction of $\Lambda(1405)$




- Two-bump structure seen
- Possible evidence for two $I=0$ poles
- PDG $\Lambda(1405)$ values fail utterly
- Calculation needed!



Strangeness Suppression of $q\bar{q}$ Creation in Exclusive Reactions

Publication: M. D. Mestayer, K. Park *et al.* (CLAS Collaboration),
Phys. Rev. Lett. **113**, 152004 (2014).

γ Y^* K^+ | $K^+\Lambda : \pi^+n : \pi^0p$ Electroproduction Ratios



Motivation:

- Quark model picture of quark-pair creation and flux-tube breaking: does it apply in the low energy exclusive limit?


Measurements:

- Ratio of processes in which only **one $q\bar{q}$ pair** is produced: an $s\bar{s}$, $d\bar{d}$, or $u\bar{u}$, respectively
- In quark model picture, ratios are proportional to the **relative production rates** of $s\bar{s}$, $d\bar{d}$, or $u\bar{u}$

Physics conclusion:

- Ratio of $s\bar{s}$ pair creation relative to $u\bar{u}$ or $d\bar{d}$ is **suppressed**; $\approx 0.2 - 0.3$
- **Consistent with high-energy results** when 100's of particles are produced






Ratio	$s\bar{s} / d\bar{d}$	$u\bar{u} / d\bar{d}$
$K^+\Lambda/\pi^+n$	0.19 ± 0.03	-
$K^+\Lambda/\pi^0p$ "a"	0.22 ± 0.07	-
$K^+\Lambda/\pi^0p$ "b"	0.28 ± 0.07	-
π^0p/π^+n	-	0.74 ± 0.18




The Future: Outlook at GlueX and CLAS12

γ Y^* K^+ Lattice QCD Predictions

- Lattice QCD now predicts rich baryon families
- Most states not identified by experiment yet




R. Edwards *et al.*, PRD 87, 054506 (2013)



Baryon Spectroscopy

- JLab at 12 GeV will surpass many Y^* thresholds
- $S = -1, -2, -3$
 - Many $\Lambda^*, \Sigma^*, \Xi^*, \Omega^*$ states remain undiscovered
- Charm threshold



(K. Moriya, priv. comm.)


γ Y^* K^+ | JLab Hall D/GlueX

- New hall, finished construction
- Commissioning in progress now
- Approved for 220 days of high statistics running





γ Y^* K^+ | Jlab Hall D/GlueX

- Real photon beam centered at 9 GeV
- Liquid hydrogen target
- Reconstruct both charged and neutral particles over large angular range
- Hermetic detector within solenoid magnetic field
- Meson & Baryon spectroscopy: search for new and exotic states






γ Y^* K^+ | GlueX Study of $\Xi^-(1820)$

- Use simulated data to study



- Final state is 5 charged particles, K^+ , K^+ , K^- , p , π^-
- Can GlueX reconstruct this?
- Reconstruction efficiency
 - 10 MeV mass resolution
 - Secondary vertex resolution: ~ 1 cm along beam line (z-direction)





JLab Hall B / CLAS12

Baseline equipment

Forward Detector (FD)

- TORUS magnet (6 coils)
- HT Cherenkov Counter
- Drift chamber system
- LT Cherenkov Counter
- Forward ToF System
- Pre-shower calorimeter
- E.M. calorimeter

Central Detector (CD)

- SOLENOID magnet
- Barrel Silicon Tracker
- Central Time-of-Flight

Beamline


- Polarized target (transv.)
- Moller polarimeter
- Photon Tagger

Upgrades to the baseline

Under construction

- MicroMegas
- Central Neutron Detector
- Forward Tagger
- RICH detector (1 sector)
- Polarized target (long.)

6/19/14




γ Y^* K^+ CLAS12: Very Strange Baryons

Study of the Ω^- and Ξ^* are among the main goals of the CLAS12 spectroscopy program:


- Ω^- discovered in 1964: after 50 years, indication on J^P from Babar and others but full determination not yet achieved
- Ξ^* spectrum still poorly known: many states missing and spin/parity undetermined


Photoproduction mechanism implies creation of three s quarks

- Models indicate $\sigma(\Omega^-) \sim 0.3-2$ nb at $E \sim 7$ GeV
- Expected production rates in CLAS12:
 - Ω^- : 90 /h
 - $\Xi^-(1690)/\Xi^-(1820)$: 0.2/0.9 k/h
- Ω^- : measurement of the cross section and investigation of production mechanisms
- Ξ^* : spin/parity determination, cross section and production mechanism, measurement of doublets mass splitting




V. E. Barnes et al., Phys. Rev. Let. 12 (1964) 204





Summary/Conclusions

- Hyperon photo- and electro-production used to pin down N^* spectrum above 1.6 GeV
- Y^* cross sections compared; $\Lambda(1405)$ "weird"
- Interference effects in $\Lambda(1405)$ line shapes demonstrated
- Direct J^P measurement for $\Lambda(1405)$ made: $\frac{1}{2}^-$
- Cross section scaling demonstrated and strangeness suppression seen
- JLab at 12 GeV with CLAS12 and GlueX will explore Y^* and meson spectra



Supplemental Slides
