



# Electromagnetic Strangeness Production at GeV Energies



Reinhard Schumacher  
**Carnegie Mellon University**

for the CLAS & GlueX Collaborations

October 9, 2014, DNP/JPS Hawaii




# Outline / Overview

- Strangeness and the  $N^*$  spectrum of states
  - Ground state hyperon photoproduction
  - Ground state hyperon electroproduction
- Dimensional scaling of  $K\Lambda$  photoproduction
- Excited  $Y^*$  cross sections measured at CLAS
  - $\Sigma^0(1385)$  ( $J^P = 3/2^+$ );  $\Lambda(1405)$  ( $J^P = 1/2^-$ );  $\Lambda(1520)$  ( $J^P = 3/2^-$ )
- The  $\Lambda(1405)$  and chiral unitary models
  - Line shapes in  $\Sigma \pi$
  - Spin & parity  $J^P$  of the  $\Lambda(1405)$
  - First **electro**-production of  $\Lambda(1405)$
- $K^*Y$  production
- Outlook at GlueX and CLAS12

$\gamma$   $Y^*$   
 $K^+$  | Jefferson Lab


- Located in Newport News, Virginia
- Ran for ~14 yrs at 6 GeV in Halls A, B, C
- Upgrading to 12 GeV, new Hall D



- Most  $Y$ ,  $Y^*$  publications from Hall B
- Upgrading as CLAS12 for 12 GeV




# CLAS Experiment




# CLAS Experiment

- Photoproduction:
  - Targets: unpolarized  $\text{LH}_2$ , polarized p, & HD-ice
  - Beams: unpolarized, circular, linear, to  $\sim 5 \text{ GeV}$
  - Reconstructed  $\text{K}^+p\pi^-(\pi^0)$  or  $\text{K}^+\pi^+\pi^-(n)$
  - $20 \times 10^9$  triggers  $\rightarrow 1.41 \times 10^6$   $\text{KY}\pi$  events in g11a
- Electroproduction:
  - $Q^2$  from  $\sim 0.5$  to  $\sim 3 (\text{GeV}/c)^2$
  - Structure functions from Rosenbluth and beam-helicity separations



# Strangeness and the $N^*$ Spectrum of States - Photoproduction



# Strangeness in $N^*$ Physics: Status

**Table 8.** Star rating suggested for baryon resonances and their decays. Ratings of the Particle Data Group are given as \*; additional stars suggested from this analysis are represented by  $\star$ ; (\*) stands for stars which should be removed.

	all	$\pi N$	$\gamma N$	$N\eta$	$\Delta K$	$\Sigma K$	$\Delta\pi$	$N\sigma$
$N(1440)_{1/2}^{+}$	****	****	****	(*)			***	***
$N(1710)_{1/2}^{+}$	***	***	***	***	***	***	*(*)	
$N(1880)_{1/2}^{+}$	**	*	*		**	*		
$N(1535)_{1/2}^{-}$	****	****	****	****			*	
$N(1650)_{1/2}^{-}$	****	****	***	***	***	**	**(*)	
$N(1895)_{1/2}^{-}$	**	*	**	**	**	*		
$N(1720)_{3/2}^{+}$	****	****	****	****	**	**	***	
$N(1900)_{3/2}^{+}$	***	**	***	**	***	***	**	
$N(1520)_{3/2}^{-}$	****	****	****	***			****	
$N(1700)_{3/2}^{-}$	**	**	**	*	*(*)	*	***	
$N(1875)_{3/2}^{-}$	***	*	***		***	***		***
$N(2150)_{3/2}^{-}$	**	**	**		**		**	
$N(1680)_{3/2}^{+}$	****	****	****	*			**(*)	**
$N(1860)_{3/2}^{+}$	*	*	*					
$N(2000)_{3/2}^{+}$	***	*(*)	**	**	**	*		
$N(1675)_{3/2}^{-}$	****	****	****(*)	*	*		***(*)	*
$N(2060)_{3/2}^{-}$	***	**	***	*		**		
$N(1990)_{7/2}^{+}$	**	*(*)	**					
$N(2190)_{7/2}^{-}$	****	****	***		**			
$N(2220)_{9/2}^{+}$	****	****						
$N(2250)_{9/2}^{-}$	****	****						
$\Delta(1910)_{1/2}^{+}$	****	****	**		**		**	
$\Delta(1620)_{1/2}^{-}$	****	****	**				****	
$\Delta(1900)_{1/2}^{-}$	**	**	**		**		**	
$\Delta(1232)_{3/2}^{+}$	****	****	****					
$\Delta(1600)_{3/2}^{+}$	***	**	**				***	
$\Delta(1920)_{3/2}^{+}$	***	**	**		***		**	
$\Delta(1700)_{3/2}^{-}$	***	**	**				**	
$\Delta(1940)_{3/2}^{-}$	*	*	**					* from $\Delta\eta$
$\Delta(1905)_{5/2}^{+}$	****	****	****		***		**(*)	
$\Delta(1950)_{7/2}^{+}$	****	****	**		***		**	



$S_{11}$   $\rightarrow$   
 $P_{13}$   $\rightarrow$   
 $D_{13}$   $\rightarrow$   
 $G_{17}$   $\rightarrow$

- What role has e.m. strangeness physics in unraveling  $N^*$  and  $\Delta$  properties?
- Worldwide effort to determine resonance poles, branching fractions, helicity couplings, etc.
- Bottom line: "Stars" and resonances added to world database

$\gamma$   $Y^*$   
 $K^+$

# Define the Spin Observables

(for NO nucleon polarization)



$$\frac{d\sigma}{d\Omega} = \sigma_0 \left\{ \begin{array}{l} 1 - P_{\bar{\gamma}} \Sigma \cos 2\phi \\ - \alpha \cos \theta_{x'} \sin 2\phi P_{\bar{\gamma}} O_{x'} - \alpha \cos \theta_{x'} P_{\odot} C_{x'} \\ - \alpha \cos \theta_{z'} \sin 2\phi P_{\bar{\gamma}} O_{z'} - \alpha \cos \theta_{z'} P_{\odot} C_{z'} \\ + \alpha \cos \theta_{y'} P - \alpha \cos \theta_{y'} P_{\bar{\gamma}} T \cos 2\phi \end{array} \right\}$$


# $\gamma$ $Y^*$ $K^+$ Observables in Pseudoscalar Meson Photoproduction

4 Complex amplitudes: **16** real polarization observables.

Complete measurement from **8** carefully chosen observables.

$nN$  has large cross section

but in  $KY$  recoil is **self-analysing**



$nN$		Symbol	Transversity representation	Experiment required	Type	$KY$		
recoil	target					$\gamma$	$\gamma$	target
		$d\sigma/dt$	$ b_1 ^2 +  b_2 ^2 +  b_3 ^2 +  b_4 ^2$	$\{-; -; -\}$	$S$			
		$\Sigma d\sigma/dt$	$ b_1 ^2 +  b_2 ^2 -  b_3 ^2 -  b_4 ^2$	$\{L(\frac{1}{2}\pi, 0); -; -\}$				
		$Td\sigma/dt$	$ b_1 ^2 -  b_2 ^2 -  b_3 ^2 +  b_4 ^2$	$\{-; y; -\}$				
		$Pd\sigma/dt$	$ b_1 ^2 -  b_2 ^2 +  b_3 ^2 -  b_4 ^2$	$\{-; -; y\}$				
		$Gd\sigma/dt$	$2 \text{Im}(b_1 b_3^* + b_2 b_4^*)$	$\{L(\pm\frac{1}{4}\pi); z; -\}$	$BT$			
		$Hd\sigma/dt$	$-2 \text{Re}(b_1 b_3^* - b_2 b_4^*)$	$\{L(\pm\frac{1}{4}\pi); x; -\}$				
		$Ed\sigma/dt$	$-2 \text{Re}(b_1 b_3^* + b_2 b_4^*)$	$\{C; z; -\}$				
		$Fd\sigma/dt$	$2 \text{Im}(b_1 b_3^* - b_2 b_4^*)$	$\{C; x; -\}$				
		$O_x d\sigma/dt$	$-2 \text{Re}(b_1 b_4^* - b_2 b_3^*)$	$\{L(\pm\frac{1}{4}\pi); -; x'\}$	$BR$			
		$O_z d\sigma/dt$	$-2 \text{Im}(b_1 b_4^* + b_2 b_3^*)$	$\{L(\pm\frac{1}{4}\pi); -; z'\}$				
		$C_x d\sigma/dt$	$2 \text{Im}(b_1 b_4^* - b_2 b_3^*)$	$\{C; -; x'\}$				
		$C_z d\sigma/dt$	$-2 \text{Re}(b_1 b_4^* + b_2 b_3^*)$	$\{C; -; z'\}$				
		$T_x d\sigma/dt$	$2 \text{Re}(b_1 b_2^* - b_3 b_4^*)$	$\{-; x; x'\}$	$TR$			
		$T_z d\sigma/dt$	$2 \text{Im}(b_1 b_2^* - b_3 b_4^*)$	$\{-; x; z'\}$				
		$L_x d\sigma/dt$	$2 \text{Im}(b_1 b_2^* + b_3 b_4^*)$	$\{-; z; x'\}$				
		$L_z d\sigma/dt$	$2 \text{Re}(b_1 b_2^* + b_3 b_4^*)$	$\{-; z; z'\}$				

I. S. Barker, A. Donnachie, J. K. Storrow, Nucl. Phys. B95 347 (1975).

circ polarized photons  
 linearly polarized photons

longitudinally polarized target  
 transversely polarized target

**Complete, and over-determined**


# $\gamma$ $Y^*$ $K^+$ Theory: Bonn Gatchina Model

(Just one of several models on the market)

- Coupled channels (K-matrix) framework
  - Input: from  $\pi N$ ,  $K N$  elastic;  $\gamma N$ ,  $\pi N$  inelastic to  $\pi^{\pm 0} N$ ,  $\eta N$ ,  $\eta' N$ ,  $K^{\pm 0} Y$ ,  $\pi \pi N$ 
    - Use ALL experimental channels, including the strangeness channels & spin observables
  - Partial Wave Analysis
    - First extract each  $J^P$  wave
    - Fit  $N^*$  and  $\Delta$  resonance pole parameters


## Short list of References:

- A. Sarantsev, V. Nikonov, A. Anisovich, E. Klempt, U. Thoma; Eur. Phys. J. A **25**, 441 (2005)  
A.V. Anisovich *et al.*, Eur. Phys J. A **25** 427 (2005); Eur. Phys J. A **24**, 111 (2005);  
V. A. Nikonov *et al.*, Phys Lett. B **662**, 246 (2008).  
A. Anisovich, E. Klempt, V. Nikonov, A. Sarantsev, U. Thoma; Eur. Phys. J. A **47**, 153 (2011).







# $\gamma p \rightarrow K^+ \Lambda$ : cross section

R. Bradford *et al.*, Phys.Rev. C **73**, 035202 (2006)



- Forward peaking indicates t-channel processes at high  $W$
- Angular dependence at lower  $W$  consistent with s- and u-channel processes.

# $\gamma Y^* K^+$ | $\gamma p \rightarrow K^+ \Lambda$ : recoil polarization $P$



- Kaon-MAID model (green)
  - F.X. Lee *et al.*, Nucl. Phys. **A695**, 237 (2001).
  - Single-channel BW resonance fits
  - No longer up-to-date
- Bonn-Gatchina model (blue)
  - Multi-channel, unitary, BW resonance fit
  - Large suite of  $N^*$  contributions
  - Was not predictive for recoil polarization

A.V. Sarantsev *et al.*, Eur. Phys. J., A **25**, 441 (2005).

# $\gamma p \rightarrow K^+ \Lambda$ : beam asymmetry $\Sigma$




Fig. 14. Angular distributions of the beam asymmetries  $\Sigma$  for  $\gamma p \rightarrow K^+ \Lambda$  and  $\gamma$ -ray energies ranging from to 1500 MeV. Data are compared with the new solutions of the BCC (solid line), SAPCC (dashed line) and GRP models.




FIG. 3. Beam polarization asymmetries for the  $p(\vec{\gamma}, K^+) \Lambda$  (left) and  $p(\vec{\gamma}, K^+) \Sigma^0$  (right) reactions as a function of  $\cos(\theta_{K^+}^{cm})$  for different photon-energy bins. The error bars are experimental.

$$\frac{d\sigma}{d\Omega_{K^+}} = \frac{d\sigma}{d\Omega_{K^+}} \Big|_{unpol.} \{1 + \Sigma P_\gamma \cos 2\phi\}$$

GRAAL threshold range,  $E_\gamma < 1.5 \text{ GeV}$


LEPS  $1.5 < E_\gamma < 2.4 \text{ GeV}$

The trends are consistent:  $\Sigma$  is smooth and featureless at all energies and angles.

LEPS

R. G. T. Zegers *et al.* (LEPS) Phys. Rev. Lett. **91**, 092001 (2003).


A. Lleres *et al.* (GRAAL) Eur. Phys. J. A **31**, 79 (2007). Carnegie Mellon University



# $\gamma p \rightarrow K^+ \Lambda$ : beam asymmetry $\Sigma$

$$\frac{d\sigma}{d\Omega_{K^+}} = \frac{d\sigma}{d\Omega_{K^+}} \Big|_{unpol.} \{1 + \Sigma P_\gamma \cos 2\phi\}$$


Observable: Beam Asymmetry,  $\Sigma$



Bonn-Gatchina model is not predictive in newly-measured kinematics





# $\gamma$ $\text{K}^+$ $\text{Y}^*$ | $\gamma p \rightarrow \text{K}^+ \Lambda$ : target asymmetry T

$$\gamma + p \rightarrow \text{K}^+ \Lambda$$



Bonn-Gatchina model is not predictive in newly-measured kinematics

# $\gamma$ $\Upsilon^*$ $K^+$ $\gamma p \rightarrow K^+ \Lambda$ : helicity asymmetry E



preliminary

$\gamma$   $Y^*$   $K^+$   $\vec{\gamma}p \rightarrow K^+ \vec{\Lambda}$  Beam-Recoil  $O_x$  and  $O_z$






Fig. 9. Angular distributions of the beam-recoil observable  $O_x$  for photon energies  $E_\gamma$  ranging from 980 MeV to 1466 MeV. Error bars represent the quadratic sum of statistical and systematic errors. Data are compared with the predictions of the BG (solid line) and RPR (dashed line) models.

GRAAL data: fair agreement with BoGa and RPR models

$\gamma$   $\gamma^*$   $K^+$   $\vec{\gamma}p \rightarrow K^+ \vec{\Lambda}$  Beam-Recoil  $O_x$  and  $O_z$

$\gamma + p \rightarrow K^+ \Lambda$


Observable: Beam-Recoil,  $O_z$



$O_x$


The Bonn-Gatchina model is (again) not predictive in newly-measured kinematics

Observable: Beam-Recoil,  $O_x$




$O_z$

# $\gamma$ $Y^*$ $K^+$ $\vec{\gamma}p \rightarrow K^+ \vec{\Lambda}$ Beam-Recoil $C_x$ and $C_z$




$C_x C_z$  without  $N^*(1900)P_{13}$



$C_x C_z$  with  $N^*(1900)P_{13}$


- Nikanov *et al.*'s refit of Bonn-Gatchina coupled-channel isobar model
- mix includes:  $S_{11}$ -wave,  $P_{13}(1720)$ ,  $P_{13}(1900)$ ,  $P_{11}(1840)$
- $K^+\Sigma^0$  cross sections also better described with  $P_{13}(1900)$



R. Bradford *et al.*, (CLAS Collaboration) Phys. Rev. C **75**, 035205 (2007).


V. A. Nikanov *et al.*, Phys Lett. B **662**, 246 (2008).

see also: A.V. Anisovich *et al.*, Eur. Phys J. A **25** 427 (2005).




# Lots more could be said...

- Omit results for  $\Sigma$  photoproduction
- Omit discussion of reactions on the neutron (deuteron), which tease isospin dependence apart. (Cf. talk by T. Kageya, KC11)
- Overall goal: measure enough observables for "complete" determination of amplitudes  $\Rightarrow$  extract  $N^*$  and  $\Delta$  content



# Seeking New $S=0$ Baryons via Mesons off the Proton: published, acquired, FroST(g9b)


	$\sigma$	$\Sigma$	$T$	$P$	$E$	$F$	$G$	$H$	$T_x$	$T_z$	$L_x$	$L_z$	$O_x$	$O_z$	$C_x$	$C_z$	CLAS run Period
$p\pi^0$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g8, g9
$n\pi^+$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g8, g9
$p\eta$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g11, g8, g9
$p\eta'$	✓	✓	✓	✓	✓	✓	✓	✓									g1, g11, g8, g9
$p\omega$	✓	✓	✓	✓	✓	✓	✓	✓									g11, g8, g9
$K^+\Lambda$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	g1, g8, g11
$K^+\Sigma^0$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	g1, g8, g11
$K^{0*}\Sigma^+$	✓										✓	✓			✓	✓	g1, g8, g11



# Seeking New S=0 Baryons via Mesons off the Neutron: published, acquired, HD-ice

	$\sigma$	$\Sigma$	$T$	$P$	$E$	$F$	$G$	$H$	$T_x$	$T_z$	$L_x$	$L_z$	$O_x$	$O_z$	$C_x$	$C_z$	CLAS run Period
$p\pi^-$	✓	✓	✓		✓	✓	✓	✓									g2, g10, g13, g14
$p\rho^-$	✓	✓	✓		✓	✓	✓	✓									g2, g10, g13, g14
$K^0\Lambda$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	g13, g14
$K^0\Sigma^0$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	g13, g14
$K^+\Sigma^-$	✓	✓	✓		✓	✓	✓	✓									g10, g13, g14
$K^{0*}\Sigma^0$	✓	✓															g10, g13

The combination of all of these measurements on proton and neutron targets represents an extremely powerful tool in the search for new baryon states.




# Ground States - Electroproduction

(cf. D. Carman KC6)

# Structure Functions

For unpolarized target & polarized  $e^-$  beam:



$$\frac{d^4\sigma}{dQ^2 dW d\Omega_K} = \Gamma(Q^2, W) \times \frac{d\sigma}{d\Omega_K}(Q^2, W, \Theta_K, \varepsilon, \phi)$$

Virtual photon flux

Meson cross section

Transverse

Transverse-transverse interference

Helicity structure


$$\frac{d\sigma}{d\Omega_K} = \sigma_T + \varepsilon_L \sigma_L + \varepsilon \sigma_{TT} \cos(2\phi) + \sqrt{2\varepsilon_L(\varepsilon+1)} \sigma_{LT} \cos(\phi) + h\sqrt{2\varepsilon_L(1-\varepsilon)} \sigma_{LT'}$$

$\sigma_u$   
"Unseparated"

Longitudinal (sensitive to  $J=0^\pm$  exchange in t-channel: kaons, diquarks)


Transverse-longitudinal interference

# $\gamma Y^* | K^+ \Lambda$ Structure Functions




$E = 5.5 \text{ GeV}$ ,  $W: \text{thr} - 2.6 \text{ GeV}$ ,  $Q^2 = 1.80, 2.60, 3.45 \text{ GeV}^2$  [Carman *et al.*, PR C **87**, 025204 (2013)]

# $\gamma Y^* | K^+ \Sigma^0$ Structure Functions



$E = 5.5 \text{ GeV}$ ,  $W: \text{thr} - 2.6 \text{ GeV}$ ,  $Q^2 = 1.80, 2.60, 3.45 \text{ GeV}^2$  [Carman *et al.*, PR C **87**, 025204 (2013)]

# $\gamma Y^* K^+$ Recoil Polarization $\vec{e}p \rightarrow e'K^+\Lambda$




$\langle Q^2 \rangle \sim 1.9 \text{ GeV}^2$

[Gabrielyan *et al.*, PR C **90**, 035202 (2014)]


*Kaon-Maid*    *Maxwell*    *RPR-2007*

*RPR-2011 (solid-full, dash-NR)*



[McCracken *et al.*, PR C **81**, 025201 (2010)]

# $\gamma Y^* K^+ |$ Transfer Polarization $\vec{e}p \rightarrow e'K^+\vec{\Lambda}$



5.754 GeV  
Summed over  $Q^2, \Phi$


- Data not included in fits
- Rule out  $P_{11}(1900)$  assignment
- $D_{13}(1900)$  not ruled out via  $P'$  data but with S.F. data

*Isobar Model - Mart*  
*Regge Model - GLV*  
*RPR w  $P_{11}(1900)$  - Ghent*  
*RPR w  $D_{13}(1900)$  - Ghent*

RPR background +  $S_{11}(1650)$ ,  
 $P_{11}(1710)$ ,  $P_{13}(1720)$ ,  $P_{13}(1900)$

[Carman *et al.*, PRC **79**, 065205 (2009)]

# $\gamma Y^* K^+$ | L/T Separation



[Ambrozewicz *et al.*, PR C **75**, 045203 (2007)]


# $\gamma$ $Y^*$ $K^+$ CLAS $e p$ Data Set Overview

#	Run	$E_b$ (GeV)	Trig.
1	e1c	2.567	900
2	e1c	4.056	370
3	e1c	4.247	620
4	e1c	4.462	420
5	e1d	4.817	300
6	e1-6	5.754	4500
7	e1f	5.499	5000
8	e1g	3.178	2500

- $K^+\Lambda$  recoil polarization
  - $W=1.6-2.7$  GeV,  $\langle Q^2 \rangle = 1.9$  GeV<sup>2</sup>  
[Gabrielyan *et al.*, PR C **90**, 035202 (2014)]

## Publications:

- $K^+\Lambda$  beam-recoil pol. transfer
  - $W=1.6-2.15$  GeV,  $Q^2=0.3 - 1.5$  GeV<sup>2</sup>  
[Carman *et al.*, PRL **90**, 131804 (2003)]
- $K^+\Lambda$   $\sigma_L/\sigma_T$  ratio from pol. transfer data
  - $W=1.72-1.98$  GeV,  $Q^2 \sim 0.7$  GeV<sup>2</sup>  
[Raue & Carman, PR C **71**, 065209 (2005)]
- $K^+\Lambda$ ,  $K^+\Sigma^0$  separated structure functions
  - $W=thr-2.4$  GeV,  $Q^2=0.5-2.8$  GeV<sup>2</sup>
  - $\sigma_U, \sigma_{LT}, \sigma_{TT}, \sigma_L, \sigma_T$  -  $K^+\Lambda, K^+\Sigma^0$   
[Ambrozewicz *et al.*, PR C **75**, 045203 (2007)]
  - $W=thr-2.6$  GeV,  $Q^2=1.4-3.9$  GeV<sup>2</sup>
  - $\sigma_U, \sigma_{LT}, \sigma_{TT}, \sigma_{LT}$  -  $K^+\Lambda, K^+\Sigma^0$   
[Carman *et al.*, PRC **87**, 025204 (2013)]
- $K^+\Lambda$  fifth structure function  $\sigma_{LT}$ 
  - $W=1.6-2.1$  GeV,  $Q^2=0.65, 1.0$  GeV<sup>2</sup>  
[Nasseripour *et al.*, PR C **77**, 065208 (2008)]
- $K^+\Lambda, K^+\Sigma^0$  beam-recoil pol. transfer
  - $W=thr-2.6$  GeV,  $Q^2=1.6-2.6$  GeV<sup>2</sup>  
[Carman *et al.*, PR C **79**, 065205 (2009)]




# Dimensional Scaling of $K\Lambda$


Publication: **Scaling and Resonances in Elementary  $K^+\Lambda$  Photo-production**, R.A.Sch. and M.M. Sargsian Phys.Rev.C**83** 025207 (2011).

# Constituent-Counting Scaling

$$\frac{d\sigma}{dt} = f\left(\frac{t}{s}\right) s^{2-n}$$




$n=9$





$n=10$

- Constituent counting rules for exclusive scattering
- Valid for  $s \rightarrow \infty$  and  $t/s$  fixed
  - $t/s \sim \cos(\theta_{\text{cm}})$  as  $s \rightarrow \infty$
- $n$  = number of point-like constituents
- Follows from pQCD... but also other models
- Does it work for  $K\Lambda$  ?




# Resonance Fit to Cross Section






# N\* Baryons: Seen & "Missing"




- Relativized CQM
  - Classify oscillator-model states by I, J, P
- Consistent with observation of a "missing" N\* state in K $\Lambda$
- PDG2013 now lists the "\*\*\*" N(2150) 3/2<sup>-</sup> D<sub>13</sub>




# Excited $Y^*$ Cross Sections

Publication: **Differential Photoproduction Cross Sections of  $\Sigma^0(1385)$ ,  $\Lambda(1405)$  and  $\Lambda(1520)$** , K. Moriya *et al.* (CLAS Collaboration), *Phys. Rev. C* **88**, 045201 (2013).




# Detect $K^+p\pi^-(\pi^0)$ or $K^+\pi^+\pi^-(n)$

counts/5 MeV  $MM(\gamma, K^+)$



# $\gamma$ $Y^*$ $K^+$ Differential $\Sigma^0(1385)$ Cross Section



- $\gamma + p \rightarrow K^+ + \Sigma^0(1385)$

- Experiment: see  $t$ -channel-like forward peaking &  $u$ -channel backward rise


- Agreement with LEPS

- Theory by Oh et al.<sup>1</sup>: contact term dominant; included four high-mass  $N^*$  and  $\Delta$  resonances

- Prediction was fitted to preliminary CLAS total cross section (years ago)

1. Y. Oh, C. M. Ko, K. Nakayama, Phys. Rev. **C 77**, 045204 (2008)

# $\gamma$ $Y^*$ $K^+$ Differential $\Lambda(1520)$ Cross Section



■  $\gamma + p \rightarrow K^+ + \Lambda(1520)$

■ Experiment: see  $t$ -channel-like forward peaking &  $u$ -channel backward rise

■ Agreement with LEPS<sup>1,2</sup>


■ Theories:

■ Nam & Kao<sup>3</sup>: contact term dominant; no  $K^*$  or  $u$ -channel exchanges

■ He & Chen<sup>4</sup>:  $K^*$  and  $N(2080)D_{13}$   $J^P=3/2^-$  added


1. H. Kohri et al. (LEPS) Phys Rev Lett **104**, 172001 (2010)
2. N. Muramatsu et al. (LEPS) Phys Rev **103**, 012001 (2009)
3. S.I. Nam & C.W. Kao, Phys. Rev. **C 81**, 055206 (2010)
4. J. He & X.R. Chen, Phys. Rev. **C 86**, 035204 (2012)

# $\gamma$ $Y^*$ $K^+$ Differential $\Lambda(1405)$ Cross Section



- $\gamma + p \rightarrow K^+ + \Lambda(1405)$
- Experiment: first-ever measurements
- High  $W$ : See  $t$ -channel-like forward peaking &  $u$ -channel backward rise at high  $W$
- Low  $W$ : See strong isospin dependence
  - Charge channels differ
  - WHY?!?
- Channels merge together at high  $W$

# $\gamma$ $Y^*$ $K^+$ Differential $\Lambda(1405)$ Cross Section



- $\gamma + p \rightarrow K^+ + \Lambda(1405)$

- Sum three  $\Sigma\pi$  decay modes  $\rightarrow$  "net" differential cross section

- Mixed agreement with LEPS data<sup>1</sup>

- Theories:

- Nam et al.<sup>2</sup>:  $s$ -channel Born term dominant ;  $K^*$  exchange for 3 values of  $g_{K^*\Lambda^*}$


- Williams, Ji, Cotanch<sup>3</sup>: crossing and duality constraints; no  $N^*$ , estimated  $g_{K\Lambda^*}$

1. M. Niyama et al. (LEPS) Phys Rev **C78**, 035202 (2008)

2. S.I. Nam et al., J. Kor. Phys. Soc. **59**, 2676 (2011)

3. R. Williams et al., Phys. Rev. **C43**, 452 (1991)






# $\Lambda(1405)$ and Chiral Unitary Models

Publications: Measurement of the  $\Sigma\pi$  Photo-production Line Shapes Near the  $\Lambda(1405)$ , K. Moriya *et al.* (CLAS Collaboration), Phys. Rev. C **87**, 035206 (2013);

Isospin Decomposition of the Photoproduced  $\Sigma\pi$  System near the  $\Lambda(1405)$ , R. A. Sch. & K. Moriya, Nucl. Phys A **914**, 51 (2013) .



# Chiral Unitary Models






Fig. 1. Trajectories of the poles in the scattering amplitudes obtained by changing the SU(3) breaking parameter  $x$  gradually. At the SU(3) symmetric limit ( $x = 0$ ), only two poles appear, one is for the singlet and the other for the octets. The symbols correspond to the step size  $\delta x = 0.1$ .

- SU(3) baryons irreps  $1+8_s+8_a$  combine with 0- Goldstone bosons to generate:
- Two octets and a singlet of  $\frac{1}{2}^-$  baryons generated dynamically in SU(3) limit
- SU(3) breaking leads to two  $S=-1$   $I=0$  poles near 1405 MeV
  - $\sim 1420$  mostly  $\bar{K}N$
  - $\sim 1390$  mostly  $\pi\Sigma$
- Possible weak  $I=1$  pole also predicted

D. Jido, J.A. Oller, E. Oset, A. Ramos, U-G Meissner Nucl. Phys. A **725**, 181 (2003)  
 J.A. Oller, U.-G. Meissner Phys. Lett B **500**, 263 (2001).




# Example at $W=2.30$ GeV



**I=1 contribution**

# Isospin Interference



Final  $\Sigma\pi$  state

$$\left. \begin{array}{l} |I, I_3\rangle = \\ |0,0\rangle, |1,0\rangle \end{array} \right\}$$


$$|t_I|^2 \equiv |\langle I, 0 | \hat{T}^{(I)} | \gamma p \rangle|^2$$

Three charge combinations:

$$|T_{\pi-\Sigma^+}|^2 = \frac{1}{3}|t_0|^2 + \frac{1}{2}|t_1|^2 - \frac{2}{\sqrt{6}}|t_0||t_1|\cos\phi_{01},$$

$$|T_{\pi^0\Sigma^0}|^2 = \frac{1}{3}|t_0|^2,$$


$$|T_{\pi^+\Sigma^-}|^2 = \frac{1}{3}|t_0|^2 + \frac{1}{2}|t_1|^2 + \frac{2}{\sqrt{6}}|t_0||t_1|\cos\phi_{01}.$$



# What "is" the I=1 piece?


- I=1 resonance? I=1 continuum amplitude?
- L. Roca and E. Oset paper<sup>1</sup>
  - Possible I=1 resonance in vicinity of  $N\bar{K}$  threshold
- B.-S. Zou papers<sup>2</sup>
  - $\Sigma\left(\frac{1}{2}\right)^-$  is a  $|[ud][us]\bar{s}\rangle$  state: part of a new nonet
- No interference seen in  $\Lambda(1520)$  mass range: therefore it's not a continuum amplitude
- More investigation needed !

1. L. Roca, E. Oset "On the isospin 0 and 1 resonances from  $\pi\Sigma$  photoproduction data" Phys. Rev. C **88** 055206 (2013).
2. Bing-Song Zou "Five-quark components in baryons", Nucl Phys A 835 199 (2010).





# "Best" model calculation

- L. Roca and E. Oset ← best job so far
  - Possible I=1 resonance in vicinity of  $N\bar{K}$  threshold





1. L. Roca, E. Oset "On the isospin 0 and 1 resonances from  $\pi\Sigma$  photoproduction data" Phys. Rev. C **88** 055206 (2013).

$\gamma$




Line Shapes



Cross Sections

- This effect is NOT seen for the  $\Lambda(1520)$
- No model calculation has computed cross section and line shapes together.



# Spin and Parity of $\Lambda(1405)$

Publication: **Spin and Parity of the  $\Lambda(1405)$  Baryon**, K. Moriya *et al.*  
(CLAS Collaboration), Phys. Rev. Lett. **112**, 082004 (2014).

# $\gamma$ $Y^*$ $K^+$ Parity and Spin of $\Lambda(1405)$

- How does one measure these things?
  - Find a reaction wherein  $\Lambda^*$  is created polarized
  - Decay angular distribution to  $\Sigma \pi$  relates to  $J$ 
    - $J = 1/2$  : flat distribution is the best possible evidence
    - $J = 3/2$  : "smile or frown" distribution, where  $p$  is the  $m = \pm 3/2$  fraction

$$I(\theta_Y) \propto 1 + \frac{3(1-2p)}{2p+1} \cos^2 \theta_Y$$


- Parity given by polarization transfer to daughter
- No model dependence: pure kinematics

$\gamma$   $Y^*$   
 $K^+$

# S-wave, P-wave Scenarios


L=0 (s-wave)

$$\vec{Q} = \vec{P}$$




L=1 (p-wave)


$$\vec{Q} = -\vec{P} + 2(\vec{P} \cdot \hat{q})\hat{q}$$



$\Lambda(1405) \rightarrow \Sigma \pi$




# Parity and Spin of $\Lambda(1405)$



$J^P = 1/2^-$  confirms quark model expectation

- Polarization axis is along  $\hat{z} = \hat{\gamma} \times \hat{K}$
- Used  $W=2.55$  to  $2.85$  GeV,  $\cos \theta_K^{c.m.} > 0.6$
- Decay  $\Lambda(1405) \rightarrow \Sigma^+ \pi^-$  is isotropic ( $p = 0.5$ ), so  $J \rightarrow 1/2$
- Weak decay asymmetry for  $\Sigma^+$  is  $\alpha = -0.98$  (big!)
- Decay is s-wave,  $\Rightarrow P = \text{"negative"}$



and  $\Lambda(1405)$  is produced  $\sim +45\%$  polarized



# $\Lambda(1405)$ Electroproduction




Publication: **First Observation of the  $\Lambda(1405)$  Line Shape in Electroproduction**, H. Lu *et al.* (CLAS Collaboration), Phys. Rev. C **88**, 045202 (2013).

# $\gamma$ $Y^*$ $K^+$ | Electroproduction of $\Lambda(1405)$




- Probe the pole "structure" for  $Q^2 > 0$  via electromagnetic form factors
- Theory: e.m. form factors computed;  $\Lambda(1405)$  is "larger" than the neutron
- Experiment: hard to isolate pure e.m.  $\gamma\Lambda^*\Lambda$  vertex
- In CLAS  $e p \rightarrow e' K^+ p \pi^- (\pi^0)$ , four particles detected
- CLAS acceptance:
  - $1.0 < Q^2 < 3.0 \text{ GeV}^2$ ;
  - $1.5 < W < 3.5 \text{ GeV}$ .

# $\gamma$ $Y^*$ $K^+$ Electroproduction of $\Lambda(1405)$




- Two-bump structure seen
- Possible evidence for two  $I=0$  poles
- PDG  $\Lambda(1405)$  values fail utterly
- Calculation needed!



# $K^*$ Production

Publication: Cross Sections for the  $\gamma p \rightarrow K^{*+} \Lambda$  and  $\gamma p \rightarrow K^{*+} \Sigma^0$  Reactions, W. Tang *et al.* (CLAS) Phys. Rev. C **87**, 065204 (2013).


$\gamma$   $Y^*$   $K^+$  |  $K^{*+}\Lambda, K^{*+0}\Sigma^{0+}$  photoproduction



- $N^*$  search with coupling to  $K^*Y$
- Search for  $\kappa$ -meson interaction

# $\gamma$ $Y^*$ $K^+$ | $K^{*+}\Lambda$ , $K^{*+0}\Sigma^{0+}$ photoproduction

## Comparison with theory



Cyan: Oh and Kim (O-K)  
Isobar Model

Blue: Kim, Nam, Oh, Kim (KNOK)  
Regge Model

Dotted curves include additional  
s-channel  $N^*$  with  $M < 2.2$  GeV  
and  $L < 3$ .

Clearly, the currently  
available theoretical  
models cannot  
reproduce the data. This  
suggests that higher-  
mass and higher-L  
resonances are needed.

- Models include known high-mass resonances

# $\gamma$ $Y^*$ $K^+$ | $K^{*+}\Lambda$ , $K^{*+0}\Sigma^{0+}$ photoproduction





FIG. 12. (Color online) Total cross section ratio of the reactions  $\gamma p \rightarrow K^{*0}\Sigma^+$  to  $\gamma p \rightarrow K^{*+}\Lambda$ . The ratio uses the present data in the denominator and data from Ref. [5] in the numerator. The dashed and solid curves are theoretical calculations from Oh and Kim [4] models I and II, respectively.

## ■ Suggestion of $\kappa$ -meson exchange

Solid: mostly t-channel  $\kappa$ -meson  
Dotted: very little  $\kappa$ -meson






There is scarce evidence for the strange scalar called the kappa ( $\kappa$ ), which is the octet partner of the  $a_0(980)$  and  $f_0(980)$  mesons. The CLAS data support an earlier claim by LEPS that also measured  $K^{*0}\Sigma^+$  photoproduction.




# The Future: Outlook at GlueX and CLAS12

# $\gamma$ $Y^*$ $K^+$ | Lattice QCD Predictions

- Lattice QCD now predicts rich baryon families
- Most states not identified by experiment yet




R. Edwards *et al.*, PRD 87, 054506 (2013)



# Baryon Spectroscopy

- JLab at 12 GeV will surpass many  $Y^*$  thresholds
- $S = -1, -2, -3$ 
  - Many  $\Lambda^*, \Sigma^*, \Xi^*, \Omega^*$  states remain undiscovered
- Charm threshold



(K. Moriya, priv. comm.)


# $\gamma$ $Y^*$ $K^+$ | Hall D/GlueX

- New hall, finished construction
- Commissioning NOW
- Approved for 120 PAC days of commissioning, 220 days of high statistics running




# $\gamma$ $Y^*$ $K^+$ | Hall D/GlueX

- Real photon beam centered at 9 GeV
- Liquid hydrogen target
- Reconstruct both charged and neutral particles over large angular range
- Hermetic detector within solenoid magnetic field
- Meson & Baryon spectroscopy: search for new and exotic states





# $\gamma$ $Y^*$ $K^+$ | GlueX Study of $\Xi^-(1820)$


- Use simulated data to study



- Final state is 5 charged particles,  $K^+$ ,  $K^+$ ,  $K^-$ ,  $p$ ,  $\pi^-$
- Can GlueX reconstruct this?
- Reconstruction efficiency
  - 10 MeV mass resolution
  - Secondary vertex resolution:  $\sim 1$  cm along beam line (z-direction)



(K. Moriya, priv. comm.)



# Hall B / CLAS12

(cf. L. Elouadrhiri 2WG2)

## Baseline equipment

### Forward Detector (FD)

- TORUS magnet (6 coils)
- HT Cherenkov Counter
- Drift chamber system
- LT Cherenkov Counter
- Forward ToF System
- Pre-shower calorimeter
- E.M. calorimeter

### Central Detector (CD)

- SOLENOID magnet
- Barrel Silicon Tracker
- Central Time-of-Flight

### Beamline


- Polarized target (transv.)
- Moller polarimeter
- Photon Tagger

## Upgrades to the baseline

### Under construction

- MicroMegas
- Central Neutron Detector
- Forward Tagger
- RICH detector (1 sector)
- Polarized target (long.)

6/19/14




# $\gamma$ $Y^*$ $K^+$ CLAS12: Very Strange Baryons

Study of the  $\Omega^-$  and  $\Xi^*$  are among the main goals of the CLAS12 spectroscopy program:


- $\Omega^-$  discovered in 1964: after 50 years, indication on  $J^P$  from Babar and others but full determination not yet achieved
- $\Xi^*$  spectrum still poorly known: many states missing and spin/parity undetermined

Photoproduction mechanism implies creation of three s quarks


- Models indicate  $\sigma(\Omega^-) \sim 0.3-2$  nb at  $E \sim 7$  GeV
- Expected production rates in CLAS12:
  - $\Omega^-$ : 90 /h
  - $\Xi^-(1690)/\Xi^-(1820)$ : 0.2/0.9 k/h
- $\Omega^-$ : measurement of the cross section and investigation of production mechanisms
- $\Xi^*$ : spin/parity determination, cross section and production mechanism, measurement of doublets mass splitting



V. E. Barnes et al., Phys. Rev. Let. 12 (1964) 204



(R. deVita, priv. comm.)



# Summary/Conclusions

- Hyperon photo- and electro-production used to pin down  $N^*$  spectrum above 1.6 GeV
- $Y^*$  e.m. cross sections compared
- Interference effects in  $\Lambda(1405)$  cross section(s) and line shapes demonstrated
- Direct  $J^P$  measurement for  $\Lambda(1405)$  made:  $\frac{1}{2}^-$
- JLab program at 12 GeV in CLAS12 and GlueX will explore  $Y^*$  and meson spectra